

Application of the Sea-Level Affecting Marshes Model (SLAMM 5.0) to Island Bay National Wildlife Refuge

Prepared For: Dr. Brian Czech, Conservation Biologist

U. S. Fish and Wildlife Service
National Wildlife Refuge System
Division of Natural Resources and Conservation Planning
Conservation Biology Program
4401 N. Fairfax Drive - MS 670
Arlington, VA 22203

September 9, 2008

Jonathan S. Clough, Warren Pinnacle Consulting, Inc.
PO Box 253, Warren VT, 05674
(802)-496-3476

Application of the Sea-Level Affecting Marshes Model (SLAMM 5.0) to Island Bay National Wildlife Refuge

Introduction.....	1
Model Summary	1
Sea-Level Rise Scenarios	2
Methods and Data Sources	3
Results	6
Discussion:	11
References	12
Appendix A: Contextual Results	14

Introduction

Tidal marshes are among the most susceptible ecosystems to climate change, especially accelerated sea level rise (SLR). Sea level is predicted to increase by 30 cm to 100 cm by 2100 based on the International Panel on Climate Change (IPCC) Special Report on Emissions Scenarios (SRES) (Meehl et al. 2007). Rising sea level may result in tidal marsh submergence (Moorhead and Brinson 1995) and habitat migration as salt marshes transgress landward and replace tidal freshwater and brackish marsh (Park et al. 1991).

In an effort to address the potential effects of sea level rise on United States national wildlife refuges, the U. S. Fish and Wildlife Service contracted the application of the SLAMM model for most Region 4 refuges. This analysis is designed to assist in the production of comprehensive conservation plans (CCPs) for each refuge. A CCP is a document that provides a framework for guiding refuge management decisions. All refuges are required by law to complete a CCP by 2012.

Model Summary

Changes in tidal marsh area and habitat type in response to sea-level rise were modeled using the Sea Level Affecting Marshes Model (SLAMM 5.0) that accounts for the dominant processes involved in wetland conversion and shoreline modifications during long-term sea level rise (Park et al. 1989; www.warrenpinnacle.com/prof/SLAMM).

Successive versions of the model have been used to estimate the impacts of sea level rise on the coasts of the U.S. (Titus et al., 1991; Lee, J.K., R.A. Park, and P.W. Mause. 1992; Park, R.A., J.K. Lee, and D. Canning 1993; Galbraith, H., R. Jones, R.A. Park, J.S. Clough, S. Herrod-Julius, B. Harrington, and G. Page. 2002; National Wildlife Federation et al., 2006; Glick, Clough, et al. 2007; Craft et al., 2009).

Within SLAMM, there are five primary processes that affect wetland fate under different scenarios of sea-level rise:

- **Inundation:** The rise of water levels and the salt boundary are tracked by reducing elevations of each cell as sea levels rise, thus keeping mean tide level (MTL) constant at zero. The effects on each cell are calculated based on the minimum elevation and slope of that cell.
- **Erosion:** Erosion is triggered based on a threshold of maximum fetch and the proximity of the marsh to estuarine water or open ocean. When these conditions are met, horizontal erosion occurs at a rate based on site-specific data.
- **Overwash:** Barrier islands of under 500 meters width are assumed to undergo overwash during each 25-year time-step due to storms. Beach migration and transport of sediments are calculated.
- **Saturation:** Coastal swamps and fresh marshes can migrate onto adjacent uplands as a response of the fresh water table to rising sea level close to the coast.

- **Accretion:** Sea level rise is offset by sedimentation and vertical accretion using average or site-specific values for each wetland category. Accretion rates may be spatially variable within a given model domain.

SLAMM Version 5.0 is the latest version of the SLAMM Model, developed in 2006/2007 and based on SLAMM 4.0. SLAMM 5.0 provides the following refinements:

- The capability to simulate fixed levels of sea-level rise by 2100 in case IPCC estimates of sea-level rise prove to be too conservative;
- Additional model categories such as “Inland Shore,” “Irregularly Flooded (Brackish) Marsh,” and “Tidal Swamp.”
- *Optional.* In a defined estuary, salt marsh, brackish marsh, and tidal fresh marsh can migrate based on changes in salinity, using a simple though geographically-realistic salt wedge model. This optional model was not used when creating results for Island Bay National Wildlife Refuge.

Model results presented in this report were produced using SLAMM version 5.0.1 which was released in early 2008 based on only minor refinements to the original SLAMM 5.0 model. Specifically, the accretion rates for swamps were modified based on additional literature review. For a thorough accounting of SLAMM model processes and the underlying assumptions and equations, please see the SLAMM 5.0.1 technical documentation (Clough and Park, 2008). This document is available at <http://warrenpinnacle.com/prof/SLAMM>


Sea-Level Rise Scenarios

The primary set of eustatic (global) sea level rise scenarios used within SLAMM was derived from the work of the Intergovernmental Panel on Climate Change (IPCC 2001). SLAMM 5 was run using the following IPCC and fixed-rate scenarios:

Scenario	Eustatic SLR by 2025 (cm)	Eustatic SLR by 2050 (cm)	Eustatic SLR by 2075 (cm)	Eustatic SLR by 2100 (cm)
A1B Mean	8	17	28	39
A1B Max	14	30	49	69
1 meter	13	28	48	100
1.5 meter	18	41	70	150

Recent literature (Chen et al., 2006, Monaghan et al., 2006) indicates that the eustatic rise in sea levels is progressing more rapidly than was previously assumed, perhaps due to the dynamic changes in ice flow omitted within the IPCC report’s calculations. A recent paper in the journal *Science* (Rahmstorf, 2007) suggests that, taking into account possible model error, a feasible range by 2100 might be 50 to 140 cm. To allow for flexibility when interpreting the results, SLAMM was also run assuming 1 meter, 1½ meters of eustatic sea-level rise by the year 2100. The A1B- maximum scenario was scaled up to produce these bounding scenarios (Figure 1).

Figure 1: Summary of SLR Scenarios Utilized



Methods and Data Sources

No high-resolution LIDAR data were found for Island Bay NWR so elevation data are based on the National Elevation Dataset (NED). An examination of the metadata of the NED indicates that the data were derived from a 1957 survey illustrated within the USGS topographic map shown below. The contour intervals that resulted from this survey are five feet. The process of creating a digital elevation map (DEM) from a contour map does attempt to interpolate between contour lines but there is considerable uncertainty in this process.

Figure 2: Island Bay NWR from USGS Map.



Looking at this map, there are no contours drawn in this region indicating that all land is below the five foot contour. This increases the uncertainty for projections of sea level rise for this site.

The National Wetlands Inventory for Island Bay is based on a photo date of 1999. This survey, when converted to 30 meter cells, suggests that the refuge is composed of approximately 27 acres of mangroves, and 2 acres of tidal flats. Model predictions of effects due to sea level rise are run forward from 1999 as this was the date of the wetlands survey.

The historic trend for Sea Level Rise was estimated at 2.4 mm/year based on long term trends measured at Fort Myers, Florida (NOAA station 8725520). Other long term trends measured within this vicinity include Naples FL (8725110) that has registered 2.02 mm/year.

The oceanic tide range was estimated at 0.474 meters using the closest NOAA station, Turtle Bay, FL (8725649). The map vertical datum of NAVD88 was related to mean tide level using data gathered from Bokellia, Charlotte Harbor, FL (8725541).

Parameters pertaining to non-mangrove marshes (i.e. accretion rates and erosion rates) are not relevant to this site as there were no wetlands identified based on the National Wetlands Inventory, nor are any wetlands predicted to appear. Default values are therefore used, though the model will not be sensitive to those choices. Within SLAMM, mangrove accretion is set to 7mm/year based on Cahoon et al. (1999) a study that used field measurements from Rookery Bay, FL.

Tidal flat erosion rates were set to 0.5 horizontal meters per year based on the effects of wave action. This is a default model value that has been used in previous SLAMM applications to Florida. No site-specific studies of horizontal erosion rates were located for this analysis.

Modeled U.S. Fish and Wildlife Service refuge boundaries are based on Approved Acquisition Boundaries as received from Kimberly Eldridge, lead cartographer with U.S. Fish and Wildlife Service, and are current as of June, 2008. Paul Tritaik, the Wildlife Refuge Manager for this refuge assisted in providing technical contacts.

The cell-size used for this analysis was 30 meter by 30 meter cells. However, the SLAMM model does track partial conversion of cells based on elevation and slope.

SLAMM INPUT PARAMETERS FOR ISLAND BAY


Site	Island Bay
NED Source Date (yyyy)	1957
NWI_photo_date (yyyy)	1999
Direction_OffShore (N S E W)	S
Historic_trend (mm/yr)	2.4
NAVD88_correction (MTL-NAVD88 in meters)	-0.218
Water Depth (m below MLW- N/A)	2
TideRangeOcean (meters: MHHW-MLLW)	0.474
TideRangeInland (meters)	0.474
Mean High Water Spring (m above MTL)	0.315
MHSW Inland (m above MTL)	0.315
Marsh Erosion (horz meters/year)	1.8
Swamp Erosion (horz meters/year)	1
TFlat Erosion (horz meters/year) [from 0.5]	0.5
Salt marsh vertical accretion (mm/yr) Final	3.9
Brackish March vert. accretion (mm/yr) Final	4.7
Tidal Fresh vertical accretion (mm/yr) Final	5.9
Beach/T.Flat Sedimentation Rate (mm/yr)	0.5
Frequency of Large Storms (yr/washover)	25
Use Elevation Preprocessor for Wetlands	TRUE

Results

Due to high accretion rates, within this simulation, mangroves don't start to significantly lose ground until one meter of sea level rise is projected by 2100. In that case, roughly 50% of mangrove loss is predicted by 2100. However, under the scenario of 1.5 meters of sea level rise by 2100, nearly all mangroves are lost.

Tidal flats within this defined national wildlife refuge are subject to significant loss due to inundation and erosion under even the lowest projected rates of sea level rise.

Maps of SLAMM input and output to follow will use the following legend:



Island Bay


IPCC Scenario A1B-Mean, 0.39 M SLR Eustatic by 2100

Results in Acres


	Initial	2025	2050	2075	2100
Mangrove	27.1	27.1	27.1	27.1	27.1
Tidal Flat	2.2	0.4	0.1	0.0	0.0
Estuarine Open Water	0.2	2.0	2.3	2.4	2.4
Total (incl. water)	29.6	29.6	29.6	29.6	29.6




Island Bay, Initial Condition



Island Bay, 2075, Scenario A1B Mean



Island Bay, 2025, Scenario A1B Mean



Island Bay, 2100, Scenario A1B Mean



Island Bay, 2050, Scenario A1B Mean

Island Bay

IPCC Scenario A1B-Max, 0.69 M SLR Eustatic by 2100

Results in Acres

	Initial	2025	2050	2075	2100
Mangrove	27.1	27.1	27.1	27.1	27.1
Tidal Flat	2.2	0.4	0.0	0.0	0.0
Estuarine Open Water	0.2	2.1	2.4	2.4	2.4
Total (incl. water)	29.6	29.6	29.6	29.6	29.6



Island Bay, Initial Condition



Island Bay, 2075, Scenario A1B Maximum



Island Bay, 2025, Scenario A1B Maximum



Island Bay, 2100, Scenario A1B Maximum



Island Bay, 2050, Scenario A1B Maximum

Island Bay

1 Meter Eustatic SLR by 2100

Results in Acres

	Initial	2025	2050	2075	2100
Mangrove	27.1	27.1	27.1	27.1	12.9
Tidal Flat	2.2	0.3	0.0	0.0	0.0
Estuarine Open Water	0.2	2.2	2.4	2.4	16.7
Total (incl. water)	29.6	29.6	29.6	29.6	29.6




Island Bay, Initial Condition



Island Bay, 2075, 1 meter eustatic SLR by 2100



Island Bay, 2025, 1 meter eustatic SLR by 2100



Island Bay, 2100, 1 meter eustatic SLR by 2100



Island Bay, 2050, 1 meter eustatic SLR by 2100

Island Bay

1.5 Meters Eustatic SLR by 2100

Results in Acres

	Initial	2025	2050	2075	2100
Mangrove	27.1	27.1	27.1	0.9	0.0
Tidal Flat	2.2	0.3	0.0	0.0	0.0
Estuarine Open Water	0.2	2.2	2.4	28.7	29.6
Total (incl. water)	29.6	29.6	29.6	29.6	29.6




Island Bay, Initial Condition



Island Bay, 2075, 1.5 meters eustatic SLR by 2100



Island Bay, 2025, 1.5 meters eustatic SLR by 2100



Island Bay, 2100, 1.5 meters eustatic SLR by 2100



Island Bay, 2050, 1.5 meters eustatic SLR by 2100

Discussion:

Under all scenarios run, tidal flats are predicted to be lost in this region due to inundation and erosion. These tidal flat predictions are uncertain due to the predicted erosion rates and uncertainty about the initial elevations of tidal flats. Additionally, the potential formation of tidal flats due to storms is not included in the SLAMM model processes.

According to these simulations, the primary dynamic affecting mangrove abundance at this site is the rate of mangrove accretion as compared to the rate of sea level rise. Because mangroves generally accrete at a high rate they are more resilient to sea level rise. However, once sea level rise exceeds mangrove accretion rates, all mangroves are predicted to quickly disappear. Precisely where this break-point exists is uncertain, especially because modeled mangrove accretion rates were based on regional, not site-specific measurements.

Elevation data are of relatively low quality at this site, based on five foot contour intervals. However, this uncertainty is probably less important than the uncertainty as to the rate of mangrove accretion in determining long-term mangrove viability.

The contextual maps presented below probably give a better overall sense of this region's predicted vulnerability to various scenarios of sea level rise.

References

- Cahoon, D.R., J. W. Day, Jr., and D. J. Reed, 1999. "The influence of surface and shallow subsurface soil processes on wetland elevation: A synthesis." *Current Topics in Wetland Biogeochemistry*, 3, 72-88.
- Chen, J. L., Wilson, C. R., Tapley, B. D., 2006 "Satellite Gravity Measurements Confirm Accelerated Melting of Greenland Ice Sheet" *Science* 2006 0: 1129007
- Clough, J.S. and R.A. Park, 2007, *Technical Documentation for SLAMM 5.0.1* February 2008, Jonathan S. Clough, Warren Pinnacle Consulting, Inc, Richard A. Park, Eco Modeling.
<http://warrenpinnacle.com/prof/SLAMM>
- Craft C, Clough J, Ehman J, Guo H, Joye S, Machmuller M, Park R, and Pennings S. Effects of Accelerated Sea Level Rise on Delivery of Ecosystem Services Provided by Tidal Marshes: A Simulation of the Georgia (USA) Coast. *Frontiers in Ecology and the Environment*. 2009; 7, doi:10.1890/070219
- Galbraith, H., R. Jones, R.A. Park, J.S. Clough, S. Herrod-Julius, B. Harrington, and G. Page. 2002. Global Climate Change and Sea Level Rise: Potential Losses of Intertidal Habitat for Shorebirds. *Waterbirds* 25:173-183.
- Glick, Clough, et al. *Sea-level Rise and Coastal Habitats in the Pacific Northwest An Analysis for Puget Sound, Southwestern Washington, and Northwestern Oregon* July 2007
<http://www.nwf.org/sealevelrise/pdfs/PacificNWSeaLevelRise.pdf>
- IPCC, 2001: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change* [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881pp.
- Lee, J.K., R.A. Park, and P.W. Mausel. 1992. Application of Geoprocessing and Simulation Modeling to Estimate Impacts of Sea Level Rise on the Northeast Coast of Florida. *Photogrammetric Engineering and Remote Sensing* 58:11:1579-1586.
- Meehl GA, Stocker TF, Collins WD, Friedlingstein P, Gaye AT, Gregory JM, Kitoh A, Knutti R, Murphy JM, Noda A, Raper SCB, Watterson IG, Weaver AJ and Zhao ZC. 2007. Global climate projections. Pp. 747-845. In: Solomon S, Qin, D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor, M and Miller HL, (eds.) *Climate change 2007: The physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK: Cambridge University Press.
- Monaghan, A. J. *et al*, 2006 "Insignificant Change in Antarctic Snowfall Since the International Geophysical Year" *Science* 2006 313: 827-831.
- Moorhead, KK and Brinson MM. 1995. Response of wetlands to rising sea level in the lower coastal plain of North Carolina. *Ecological Applications* 5: 261-271.


- National Wildlife Federation et al., *An Unfavorable Tide: Global Warming, Coastal Habitats and Sportfishing in Florida* 4, 6 (2006).
<http://www.targetglobalwarming.org/files/AnUnfavorableTideReport.pdf>
- Park, R.A., J.K. Lee, and D. Canning. 1993. Potential Effects of Sea Level Rise on Puget Sound Wetlands. *Geocarto International* 8(4):99-110.
- Park, R.A., M.S. Trehan, P.W. Mausel, and R.C. Howe. 1989a. The Effects of Sea Level Rise on U.S. Coastal Wetlands. In *The Potential Effects of Global Climate Change on the United States: Appendix B - Sea Level Rise*, edited by J.B. Smith and D.A. Tirpak, 1-1 to 1-55. EPA-230-05-89-052. Washington, D.C.: U.S. Environmental Protection Agency.
- Rahmstorf, Stefan 2007, "A Semi-Empirical Approach to Projecting Future Sea-Level Rise," *Science* 2007 315: 368-370.
- Reed, D.J., D.A. Bishara, D.R. Cahoon, J. Donnelly, M. Kearney, A.S. Kolker, L.L. Leonard, R.A. Orson, and J.C. Stevenson, 2008: "Site-Specific Scenarios for Wetlands Accretion in the Mid-Atlantic Region. Section 2.1" in *Background Documents Supporting Climate Change Science Program Synthesis and Assessment Product 4.1: Coastal Elevations and Sensitivity to Sea Level Rise*, J.G. Titus and E.M. Strange (eds.), EPA430R07004, Washington, DC: U.S. EPA.
http://www.epa.gov/climatechange/effects/downloads/section2_1.pdf
- Stevenson and Kearney, 2008, "Impacts of Global Climate Change and Sea-Level Rise on Tidal Wetlands" Pending chapter of manuscript by University of California Press.
- Titus, J.G., R.A. Park, S.P. Leatherman, J.R. Weggel, M.S. Greene, P.W. Mausel, M.S. Trehan, S. Brown, C. Grant, and G.W. Yohe. 1991. Greenhouse Effect and Sea Level Rise: Loss of Land and the Cost of Holding Back the Sea. *Coastal Management* 19:2:171-204.

Appendix A: Contextual Results


The SLAMM model does take into account the context of the surrounding lands or open water when calculating effects. For example, erosion rates are calculated based on the maximum fetch (wave action) which is estimated by assessing contiguous open water to a given marsh cell. Another example is that inundated dry lands will convert to marshes or ocean beach depending on their proximity to open ocean.

For this reason, an area larger than the boundaries of the USFWS refuge was modeled. These results maps are presented here with the following caveats:


- Results were closely examined (quality assurance) within USFWS refuges but not closely examined for the larger region.
- Site-specific parameters for the model were derived for USFWS refuges whenever possible and may not be regionally applicable.
- Especially in areas where dikes are present, an effort was made to assess the probable location and effects of dikes for USFWS refuges, but this effort was not made for surrounding areas.




Location of Island Bay NWR within Contextual Simulation




Island Bay Context, Initial Condition




Island Bay Context, 2025, Scenario A1B Mean




Island Bay Context, 2050, Scenario A1B Mean




Island Bay Context, 2075, Scenario A1B Mean




Island Bay Context, 2100, Scenario A1B Mean




Island Bay Context, Initial Condition




Island Bay Context, 2025, Scenario A1B Maximum




Island Bay Context, 2050, Scenario A1B Maximum




Island Bay Context, 2075, Scenario A1B Maximum




Island Bay Context, 2100, Scenario A1B Maximum




Island Bay Context, Initial Condition




Island Bay Context, 2025, 1 meter eustatic SLR by 2100




Island Bay Context, 2050, 1 meter eustatic SLR by 2100




Island Bay Context, 2075, 1 meter eustatic SLR by 2100




Island Bay Context, 2100, 1 meter eustatic SLR by 2100




Island Bay Context, Initial Condition




Island Bay Context, 2025 1.5 Meters Eustatic by 2100



Island Bay Context, 2050 1.5 Meters Eustatic by 2100



Island Bay Context, 2075 1.5 Meters Eustatic by 2100



Island Bay Context, 2100 1.5 Meters Eustatic by 2100