

# **Application of the Sea-Level Affecting Marshes Model (SLAMM 6) to Pelican Island NWR**

U. S. Fish and Wildlife Service  
National Wildlife Refuge System  
Division of Natural Resources and Conservation Planning  
Conservation Biology Program  
4401 N. Fairfax Drive - MS 670  
Arlington, VA 22203

September 9, 2010

Jonathan S. Clough & Evan C. Larson, Warren Pinnacle Consulting, Inc.  
PO Box 253, Warren VT, 05674  
(802)-496-3476

# Application of the Sea-Level Affecting Marshes Model (SLAMM 6) to Pelican Island NWR

<b>Introduction.....</b>	<b>1</b>
<b>Model Summary .....</b>	<b>1</b>
Sea Level Rise Scenarios.....	1
<b>Methods and Data Sources .....</b>	<b>4</b>
<b>Results .....</b>	<b>10</b>
<b>Discussion .....</b>	<b>41</b>
<b>References .....</b>	<b>42</b>
<b>Appendix A: Contextual Results .....</b>	<b>44</b>

## Introduction

Tidal marshes are among the most susceptible ecosystems to climate change, especially accelerated sea level rise (SLR). The Intergovernmental Panel on Climate Change (IPCC) Special Report on Emissions Scenarios (SRES) suggested that global sea level will increase by approximately 30 cm to 100 cm by 2100 (IPCC 2001). Rahmstorf (2007) suggests that this range may be too conservative and that the feasible range by 2100 is 50 to 140 cm. Rising sea levels may result in tidal marsh submergence (Moorhead and Brinson 1995) and habitat “migration” as salt marshes transgress landward and replace tidal freshwater and Irregularly Flooded marsh (Park et al. 1991).

In an effort to address the potential effects of sea level rise on United States national wildlife refuges, the U. S. Fish and Wildlife Service contracted the application of the SLAMM model for most Region 1 refuges. This analysis is designed to assist in the production of comprehensive conservation plans (CCPs) for each refuge along with other long-term management plans.

## Model Summary

Changes in tidal marsh area and habitat type in response to sea-level rise were modeled using the Sea Level Affecting Marshes Model (SLAMM 6) that accounts for the dominant processes involved in wetland conversion and shoreline modifications during long-term sea level rise (Park et al. 1989; [www.warrenpinnacle.com/prof/SLAMM](http://www.warrenpinnacle.com/prof/SLAMM)).

Successive versions of the model have been used to estimate the impacts of sea level rise on the coasts of the U.S. (Titus et al., 1991; Lee, J.K., R.A. Park, and P.W. Mause. 1992; Park, R.A., J.K. Lee, and D. Canning 1993; Galbraith, H., R. Jones, R.A. Park, J.S. Clough, S. Herrod-Julius, B. Harrington, and G. Page. 2002; National Wildlife Federation et al., 2006; Glick, Clough, et al. 2007; Craft et al., 2009).

Within SLAMM, there are five primary processes that affect wetland fate under different scenarios of sea-level rise:

- **Inundation:** The rise of water levels and the salt boundary are tracked by reducing elevations of each cell as sea levels rise, thus keeping mean tide level (MTL) constant at zero. The effects on each cell are calculated based on the minimum elevation and slope of that cell.
- **Erosion:** Erosion is triggered based on a threshold of maximum fetch and the proximity of the marsh to estuarine water or open ocean. When these conditions are met, horizontal erosion occurs at a rate based on site-specific data.
- **Overwash:** Barrier islands of under 500 meters width are assumed to undergo overwash during each specified interval for large storms. Beach migration and transport of sediments are calculated.
- **Saturation:** Coastal swamps and fresh marshes can migrate onto adjacent uplands as a response of the fresh water table to rising sea level close to the coast.

- **Accretion:** Sea level rise is offset by sedimentation and vertical accretion using average or site-specific values for each wetland category. Accretion rates may be spatially variable within a given model domain or can be specified to respond to feedbacks such as frequency of flooding.

SLAMM Version 6.0 was developed in 2008/2009 and is based on SLAMM 5. SLAMM 6.0 provides backwards compatibility to SLAMM 5, that is, SLAMM 5 results can be replicated in SLAMM 6. However, SLAMM 6 also provides several optional capabilities.

- **Accretion Feedback Component:** Feedbacks based on wetland elevation, distance to channel, and salinity may be specified. This feedback will be used in USFWS simulations, but only where adequate data exist for parameterization.
- **Salinity Model:** Multiple time-variable freshwater flows may be specified. Salinity is estimated and mapped at MLLW, MHHW, and MTL. Habitat switching may be specified as a function of salinity. This optional sub-model is not utilized in USFWS simulations.
- **Integrated Elevation Analysis:** SLAMM will summarize site-specific categorized elevation ranges for wetlands as derived from LiDAR data or other high-resolution data sets. This functionality is used in USFWS simulations to test the SLAMM conceptual model at each site. The causes of any discrepancies are then tracked down and reported on within the model application report.
- **Flexible Elevation Ranges for land categories:** If site-specific data indicate that wetland elevation ranges are outside of SLAMM defaults, a different range may be specified within the interface. In USFWS simulations, the use of values outside of SLAMM defaults is rarely utilized. If such a change is made, the change and the reason for it are fully documented within the model application reports.
- Many other graphic user interface and memory management improvements are also part of the new version including an updated *Technical Documentation*, and context sensitive help files.

For a thorough accounting of SLAMM model processes and the underlying assumptions and equations, please see the SLAMM 6.0 *Technical Documentation* (Clough, Park, Fuller, 2010). This document is available at <http://warrenpinnacle.com/prof/SLAMM>

All model results are subject to uncertainty due to limitations in input data, incomplete knowledge about factors that control the behavior of the system being modeled, and simplifications of the system (CREM 2008). Site-specific factors that increase or decrease model uncertainty may be covered in the *Discussion* section of this report.

## Sea Level Rise Scenarios


SLAMM 6 was run using scenario A1B from the Special Report on Emissions Scenarios (SRES) – mean and maximum estimates. The A1 family of scenarios assumes that the future world includes rapid economic growth, global population that peaks in mid-century and declines thereafter, and the rapid introduction of new and more efficient technologies. In particular, the A1B scenario assumes that energy sources will be balanced across all sources. Under the A1B scenario, the IPCC WGI Fourth Assessment Report (IPCC, 2007) suggests a likely range of 0.21 to 0.48 meters of sea level rise by 2090-2099 “excluding future rapid dynamical changes in ice flow.” The A1B-mean scenario

that was run as a part of this project falls near the middle of this estimated range, predicting 0.39 meters of global sea level rise by 2100. A1B-maximum predicts 0.69 meters of global SLR by 2100.

The latest literature (Chen et al., 2006, Monaghan et al., 2006) indicates that the eustatic rise in sea levels is progressing more rapidly than was previously assumed, perhaps due to the dynamic changes in ice flow omitted within the IPCC report's calculations. A recent paper in the journal *Science* (Rahmstorf, 2007) suggests that, taking into account possible model error, a feasible range by 2100 of 50 to 140 cm. This work was recently updated and the ranges were increased to 75 to 190 cm (Vermeer and Rahmstorf, 2009). Pfeffer et al. (2008) suggests that 2 meters by 2100 is at the upper end of plausible scenarios due to physical limitations on glaciological conditions. A recent US intergovernmental report states "Although no ice-sheet model is currently capable of capturing the glacier speedups in Antarctica or Greenland that have been observed over the last decade, including these processes in models will very likely show that IPCC AR4 projected sea level rises for the end of the 21st century are too low." (US Climate Change Science Program, 2008) A recent paper by Grinsted et. al. (2009) states that "sea level 2090-2099 is projected to be 0.9 to 1.3 m for the A1B scenario..." Grinsted also states that there is a "low probability" that SLR will match the lower IPCC estimates.


To allow for flexibility when interpreting the results, SLAMM was also run assuming 1 meter, 1½ meters, and 2 meters of eustatic sea-level rise by the year 2100. The A1B- maximum scenario was scaled up to produce these bounding scenarios (Figure 1).

**Figure 1: Summary of SLR Scenarios Utilized**



## Methods and Data Sources


The digital elevation map used in this simulation was derived from LiDAR data as supplied by Florida Division of Emergency Management (FDEM) via NOAA with a timestamp of 2007 (Figure 1).



**Figure 1:** Shade-relief elevation map of refuge (red).

The wetlands layer for the study area was produced by the National Wetlands Inventory and was based on a 2006 photo date (Figure 2). Converting the NWI survey into 10 meter cells indicates that the approximately 5940 acre refuge (approved acquisition boundary including water) is composed of the following categories (excluding categories below 1%).

	Estuarine Open Water	72.7%
	Mangrove	10.2%
	Undeveloped Dry Land	9.9%
	Tidal Flat	3.4%
	Regularly Flooded Marsh	2.5%



**Figure 2:** Portion of study area for Pelican Island NWR.


According to the National Wetland Inventory, there are no impounded or diked areas within Pelican Island NWR. However, according to assistant refuge manager Nick Wirwa, several regularly flooded, tidal flat and mangrove areas are impounded (Figure 3). These areas were designated as diked in SLAMM input layers before the model run. Impounded marshes are predicted to be protected from SLR of up to 2 meters within SLAMM simulations. Impounded tidal flats are not protected within SLAMM and are therefore predicted to succumb when their elevations fall below MLLW.



**Figure 3:** Impoundments within the refuge boundary (image courtesy of Nick Wirwa, USFWS).

The historic trend for sea-level rise was estimated at 2.3 mm/year using the value of the two nearest NOAA gages with SLR data (8723170, Miami Beach, FL; 8721120, Daytona Beach Shores, FL). The rate of sea level rise for this refuge has been slightly higher than the global average for the last 100 years (approximately 1.7 mm/year, IPCC 2007a).

Two tide gages were used to estimate the tide range within this site (Figure 3). Tides in the northern portion of the refuge were estimated at 0.14 meters (great-diurnal tide or GT) using the gage at Sebastian, Indian River, FL (8722029). The southern portion of the refuge was parameterized with a tide of 0.15 meters using the gage at Wabasso, Indian River, FL (8722059).



**Figure 4:** Location of NOAA tides gages used for Pelican Island NWR.

The elevation at which estuarine water is predicted to regularly inundate the land (the salt elevation) was estimated based on a frequency of inundation analysis using data from the Port of West Palm Beach, FL gage (8722588). This procedure was done to include the effects of wind tides within estimates of land inundation. The saline inundation elevation was assumed to occur where water penetrates at least once every 30 days. The derived salt elevation from West Palm Beach was then adjusted for the microtidal regime of Pelican Island. The result was an estimate that salt water regularly penetrates an elevation approximately 0.21 meters above mean tide level (MTL) at the north of the site and 0.22 meters above MTL at the south.


No accretion studies from eastern Florida were located as part of this analysis. Accretion rates in salt marshes were set to 3.9 mm/year, based on measured rates of accretion in Georgia Marshes (Craft et. al, 2009). This value is also very close to several studies measuring accretion rates on the Gulf Coast of Florida (St. Marks FL, 4.0 mm/year from Cahoon et. al. 1995, and Hendrickson, J.C. 1997; Ochlockonee River FL, 4.05 mm/year from Hendrickson, J.C. 1997).

The MTL to NAVD88 correction was derived using the NOAA gages. Across tide gages the correction ranged from -0.35 meters to -0.27 meters.

Modeled U.S. Fish and Wildlife Service refuge boundaries for Florida are based on Approved Acquisition Boundaries as published on the FWS National Wildlife Refuge Data and Metadata website. The cell-size used for this analysis was 10 meter by 10 meter cells. Note that the SLAMM model will track partial conversion of cells based on elevation and slope.

## SUMMARY OF SLAMM INPUT PARAMETERS FOR PELICAN ISLAND NWR

Parameter	Global	SubSite 1	SubSite 2
Description		North	South
NWI Photo Date (YYYY)	2006	2006	2006
DEM Date (YYYY)	2007	2007	2007
Direction Offshore [n,s,e,w]	East	West	West
Historic Trend (mm/yr)	2.3	2.3	2.3
MTL-NAVD88 (m)	-0.35	-0.27	-0.27
GT Great Diurnal Tide Range (m)	1.2	0.14	0.15
Salt Elev. (m above MTL)	0.81	0.21	0.22
Marsh Erosion (horz. m /yr)	1.8	1.8	1.8
Swamp Erosion (horz. m /yr)	1	1	1
T.Flat Erosion (horz. m /yr)	0.5	0.5	0.5
Reg. Flood Marsh Accr (mm/yr)	3.9	3.9	3.9
Irreg. Flood Marsh Accr (mm/yr)	4.7	4.7	4.7
Tidal Fresh Marsh Accr (mm/yr)	5.9	5.9	5.9
Beach Sed. Rate (mm/yr)	0.5	0.5	0.5
Freq. Overwash (years)	15	15	15
Use Elev Pre-processor [True,False]	FALSE	FALSE	FALSE



**Figure 5:** Input subsites for model application.



## Results

SLAMM predicts that Pelican Island NWR will be variably impacted depending on the SLR scenario and wetland class. Up to 84% of refuge dry land is predicted to be lost in the highest scenarios examined, but as little as 4% of dry land is lost under the lowest. Mangrove acreage is predicted to increase by up to 19% in SLR scenarios of up to 1 meter (negative loss rates shown below). However, once SLR exceeds 1 meter, mangrove losses of up to 47% are predicted.

<b>SLR by 2100 (m)</b>	<b>0.39</b>	<b>0.69</b>	<b>1</b>	<b>1.5</b>	<b>2</b>
Mangrove	-3%	-19%	-10%	42%	47%
Undeveloped Dry Land	4%	23%	40%	65%	84%
Tidal Flat	64%	79%	86%	94%	95%
Regularly Flooded Marsh	1%	2%	7%	9%	10%

**Predicted Loss Rates of Land Categories by 2100 Given Simulated Scenarios of Eustatic Sea Level Rise**

Maps of SLAMM input and output to follow will use the following legend:

Undeveloped Dry Land	
Swamp	
Tidal Swamp	
Tidal Fresh Marsh	
Riverine Tidal	
Inland Open Water	
Inland Fresh Marsh	
Cypress Swamp	
Irregularly Flooded Marsh	
Estuarine Open Water	
Regularly Flooded Marsh	
Transitional Salt Marsh	
Tidal Flat	


Application of the Sea-Level Affecting Marshes Model (SLAMM 6) to Pelican Island NWR

Pelican Bay NWR


IPCC Scenario A1B-Mean, 0.39 M SLR Eustatic by 2100

Results in Acres


		Initial	2025	2050	2075	2100
	Estuarine Open Water	4321.2	4334.8	4373.4	4428.4	4463.9
	Mangrove	608.0	607.7	610.3	615.5	625.1
	Undeveloped Dry Land	588.2	584.2	581.4	573.8	563.9
	Tidal Flat	202.3	197.5	159.1	107.7	72.9
	Regularly Flooded Marsh	149.9	149.3	149.3	148.7	148.7
	Developed Dry Land	36.4	36.4	36.4	36.4	36.4
	Inland Open Water	12.3	12.3	12.3	12.3	12.3
	Tidal Fresh Marsh	5.6	5.5	5.5	5.5	5.5
	Inland Shore	5.5	1.7	1.6	1.2	0.8
	Irregularly Flooded Marsh	5.3	5.3	5.3	5.3	5.3
	Inland Fresh Marsh	5.0	5.0	5.0	5.0	5.0
	Estuarine Beach	0.3	0.4	0.3	0.3	0.3
	<b>Total (incl. water)</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>




Pelican Island NWR, Initial Condition




Pelican Island NWR, 2025, Scenario A1B Mean



Pelican Island NWR, 2050, Scenario A1B Mean



Pelican Island NWR, 2075, Scenario A1B Mean



Pelican Island NWR, 2100, Scenario A1B Mean


Application of the Sea-Level Affecting Marshes Model (SLAMM 6) to Pelican Island NWR

Pelican Bay NWR


IPCC Scenario A1B-Max, 0.69 M SLR Eustatic by 2100

Results in Acres


		Initial	2025	2050	2075	2100
	Estuarine Open Water	4321.2	4337.1	4411.5	4461.3	4508.2
	Mangrove	608.0	608.2	614.9	633.4	724.9
	Undeveloped Dry Land	588.2	583.6	576.4	554.5	455.2
	Tidal Flat	202.3	195.3	121.9	77.4	42.7
	Regularly Flooded Marsh	149.9	149.3	149.2	148.2	146.9
	Developed Dry Land	36.4	36.4	36.4	36.4	36.3
	Inland Open Water	12.3	12.3	12.3	12.3	10.8
	Tidal Fresh Marsh	5.6	5.5	5.5	5.4	5.2
	Inland Shore	5.5	1.7	1.4	0.7	0.3
	Irregularly Flooded Marsh	5.3	5.3	5.3	5.3	5.3
	Inland Fresh Marsh	5.0	5.0	5.0	5.0	4.2
	Estuarine Beach	0.3	0.3	0.3	0.2	0.1
	<b>Total (incl. water)</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>




Pelican Island NWR, Initial Condition




Pelican Island NWR, 2025, Scenario A1B Maximum



Pelican Island NWR, 2050, Scenario A1B Maximum



Pelican Island NWR, 2075, Scenario A1B Maximum



Pelican Island NWR, 2100, Scenario A1B Maximum


Application of the Sea-Level Affecting Marshes Model (SLAMM 6) to Pelican Island NWR

Pelican Bay NWR


1 Meter Eustatic SLR by 2100

Results in Acres


		Initial	2025	2050	2075	2100
	Estuarine Open Water	4321.2	4345.4	4437.2	4508.2	4696.7
	Mangrove	608.0	608.8	620.3	696.3	670.1
	Undeveloped Dry Land	588.2	582.8	568.8	471.0	350.5
	Tidal Flat	202.3	187.2	99.4	57.0	27.4
	Regularly Flooded Marsh	149.9	149.3	148.9	146.0	139.7
	Developed Dry Land	36.4	36.4	36.4	36.3	36.0
	Inland Open Water	12.3	12.3	12.3	10.9	10.7
	Tidal Fresh Marsh	5.6	5.5	5.3	4.4	2.4
	Inland Shore	5.5	1.7	1.0	0.3	0.1
	Irregularly Flooded Marsh	5.3	5.3	5.3	5.3	5.3
	Inland Fresh Marsh	5.0	5.0	5.0	4.1	1.2
	Estuarine Beach	0.3	0.3	0.3	0.1	0.0
	<b>Total (incl. water)</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>




Pelican Island NWR, Initial Condition




Pelican Island NWR, 2025, 1 Meter



Pelican Island NWR, 2050, 1 Meter



Pelican Island NWR, 2075, 1 Meter



Pelican Island NWR, 2100, 1 Meter


Application of the Sea-Level Affecting Marshes Model (SLAMM 6) to Pelican Island NWR

Pelican Bay NWR


1.5 Meters Eustatic SLR by 2100

Results in Acres


		Initial	2025	2050	2075	2100
	Estuarine Open Water	4321.2	4373.0	4482.9	4816.3	5170.8
	Mangrove	608.0	608.9	627.5	542.6	351.0
	Undeveloped Dry Land	588.2	581.5	540.7	353.0	205.2
	Tidal Flat	202.3	161.0	77.8	36.1	12.6
	Regularly Flooded Marsh	149.9	149.2	147.2	138.1	135.9
	Developed Dry Land	36.4	36.4	36.3	36.0	32.6
	Inland Open Water	12.3	12.3	12.2	10.7	8.1
	Tidal Fresh Marsh	5.6	5.4	4.3	1.2	0.5
	Inland Shore	5.5	1.6	0.6	0.1	0.0
	Irregularly Flooded Marsh	5.3	5.3	5.3	5.3	5.3
	Inland Fresh Marsh	5.0	5.0	5.0	0.6	0.0
	Estuarine Beach	0.3	0.3	0.2	0.0	12.4
	Ocean Beach	0.0	0.0	0.0	0.0	5.7
	<b>Total (incl. water)</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>




Pelican Island NWR, Initial Condition




Pelican Island NWR, 2025, 1.5 Meters



Pelican Island NWR, 2050, 1.5 Meters



Pelican Island NWR, 2075, 1.5 Meters



Pelican Island NWR, 2100, 1.5 Meters


Application of the Sea-Level Affecting Marshes Model (SLAMM 6) to Pelican Island NWR

Pelican Bay NWR


2 Meters Eustatic SLR by 2100

Results in Acres


		Initial	2025	2050	2075	2100
	Estuarine Open Water	4321.2	4398.1	4636.4	5049.1	5318.3
	Mangrove	608.0	610.0	578.1	411.6	320.0
	Undeveloped Dry Land	588.2	578.1	460.0	260.7	93.0
	Tidal Flat	202.3	138.8	63.1	23.2	9.9
	Regularly Flooded Marsh	149.9	149.0	143.1	136.1	135.2
	Developed Dry Land	36.4	36.4	36.3	34.2	26.1
	Inland Open Water	12.3	12.3	10.8	10.7	7.5
	Tidal Fresh Marsh	5.6	5.3	2.4	0.5	0.2
	Inland Shore	5.5	1.5	0.3	0.0	0.0
	Irregularly Flooded Marsh	5.3	5.3	5.3	5.3	5.3
	Inland Fresh Marsh	5.0	5.0	4.1	0.0	0.0
	Estuarine Beach	0.3	0.3	0.1	2.5	10.2
	Open Ocean	0.0	0.0	0.0	0.0	9.9
	Ocean Beach	0.0	0.0	0.0	6.1	4.4
	<b>Total (incl. water)</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>	<b>5940.0</b>




Pelican Island NWR, Initial Condition




Pelican Island NWR, 2025, 2 Meters



Pelican Island NWR, 2050, 2 Meters



Pelican Island NWR, 2075, 2 Meters



Pelican Island NWR, 2100, 2 Meters

## Discussion

Loss of dry lands to mangrove expansion is probably the most significant prediction within this set of SLAMM simulations. Dry-land elevations at this refuge are relatively low, and given SLR scenarios of 0.69 meters and above, over one quarter of refuge dry land is predicted to be lost. Up to 84% of dry land is predicted to be lost under the worst-case scenario (two meters of eustatic SLR by 2100). High-vertical resolution LiDAR elevation data reduce model uncertainty with regards to these predictions.

As noted above, due to this inundation of dry lands, mangrove habitat is predicted to increase in acreage during early SLR scenarios. Mangroves have been shown to have a relatively high vertical accretion rate (Cahoon et al. 1999) and also can thrive over a relatively wide elevation range. However, once SLR exceeds 1 meter by 2100, mangrove habitats rapidly begin to decline. This model does not include an accounting of potential killing frosts. Model uncertainty for this application is also increased by the reliance on off-site accretion data.

Model results for tidal flats assume that this category is not protected by levees. The SLAMM dike and impoundment model does not currently protect tidal flats; an omission to be corrected in future versions of the model. At this time, tidal flat losses at this site should be assumed to be overstated or should be equivalent to tidal-flat losses if the sea-wall around the tidal flat (central in the site) is not maintained. Regularly flooded marshes within impounded areas are assumed to be protected by those seawalls against SLR of up to two meters.

## References

- Cahoon, D.R., J. W. Day, Jr., and D. J. Reed, 1999. "The influence of surface and shallow subsurface soil processes on wetland elevation: A synthesis." *Current Topics in Wetland Biogeochemistry*, 3, 72-88.
- Cahoon, D. R., D.J. Reed, and J.W. Day, Jr. 1995. Estimating shallow subsidence in microtidal salt marshes of the southeastern United States: Kaye and Barghoorn revisited. *Marine Geology* 128:1-9.
- Chen, J. L., Wilson, C. R., Tapley, B. D., 2006 "Satellite Gravity Measurements Confirm Accelerated Melting of Greenland Ice Sheet" *Science* 2006 0: 1129007
- Clark, J. S. and W. A. Patterson III. 1984. Pollen, Pb-210 and sedimentation in the intertidal environment. *Journal of Sedimentary Petrology* 54(4):1249-1263.
- Clough, J.S. Park, R.A. and R. Fuller, 2010, *SLAMM Technical Documentation, Release 6.0 beta, Draft*, January 2010, <http://warrenpinnacle.com/prof/SLAMM>
- Craft C, Clough J, Ehman J, Guo H, Joye S, Machmuller M, Park R, and Pennings S. Effects of Accelerated Sea Level Rise on Delivery of Ecosystem Services Provided by Tidal Marshes: A Simulation of the Georgia (USA) Coast. *Frontiers in Ecology and the Environment*. 2009; 7, doi:10.1890/070219
- Council for Regulatory Environmental Modeling, (CREM) 2008. *Draft guidance on the development, evaluation, and application of regulatory environmental models* P Pascual, N Stiber, E Sunderland - Washington DC: Draft, August 2008
- Erwin, RM, GM Sanders, DJ Prosser, and DR Cahoon. 2006. High tides and rising seas: potential effects on estuarine waterbirds. Pages 214-228 in: *Terrestrial Vertebrates of Tidal Marshes: Evolution, Ecology, and Conservation* (R. Greenberg, J. Maldonado, S. Droegge, and M.V. McDonald, eds.). *Studies in Avian Biology* No. 32, Cooper Ornithological Society.
- Glick, Clough, et al. *Sea-level Rise and Coastal Habitats in the Pacific Northwest An Analysis for Puget Sound, Southwestern Washington, and Northwestern Oregon* July 2007  
<http://www.nwf.org/sealevelrise/pdfs/PacificNWSeaLevelRise.pdf>
- Hendrickson, J.C. 1997. Coastal wetland response to rising sea-level: quantification of short- and long-term accretion and subsidence, northeastern Gulf of Mexico. MS thesis, Florida State University, Tallahassee, FL. USA.
- IPCC, 2001: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change* [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881pp.
- IPCC, 2007. *Climate Change 2007 - The Physical Science Basis: Contribution of Working Group I to the Fourth Assessment Report of the IPCC*. Cambridge: Cambridge University Press. ISBN 978 0521 88009-1.


- Lee, J.K., R.A. Park, and P.W. Mausel. 1992. Application of Geoprocessing and Simulation Modeling to Estimate Impacts of Sea Level Rise on the Northeast Coast of Florida. *Photogrammetric Engineering and Remote Sensing* 58:11:1579-1586.
- Monaghan, A. J. *et al*, 2006 “Insignificant Change in Antarctic Snowfall Since the International Geophysical Year” *Science* 2006 313: 827-831.
- National Wildlife Fed’n et al., *An Unfavorable Tide: Global Warming, Coastal Habitats and Sportfishing in Florida* 4, 6 (2006).  
<http://www.targetglobalwarming.org/files/AnUnfavorableTideReport.pdf>
- Park, R.A., J.K. Lee, and D. Canning. 1993. Potential Effects of Sea Level Rise on Puget Sound Wetlands. *Geocarto International* 8(4):99-110.
- Park, R.A., M.S. Trehan, P.W. Mausel, and R.C. Howe. 1989a. The Effects of Sea Level Rise on U.S. Coastal Wetlands. In *The Potential Effects of Global Climate Change on the United States: Appendix B - Sea Level Rise*, edited by J.B. Smith and D.A. Tirpak, 1-1 to 1-55. EPA-230-05-89-052. Washington, D.C.: U.S. Environmental Protection Agency.
- Park, RA, JK Lee, PW Mausel and RC Howe. 1991. Using remote sensing for modeling the impacts of sea level rise. *World Resources Review* 3:184-220.
- Pfeffer, Harper, O’Neel, 2008. Kinematic Constraints on Glacier Contributions to 21st-Century Sea-Level Rise. *Science*, Vol. 321, No. 5894. (5 September 2008), pp. 1340-134
- Rahmstorf, Stefan 2007, “A Semi-Empirical Approach to Projecting Future Sea-Level Rise,” *Science* 2007 315: 368-370.
- Reed, D.J., D.A. Bishara, D.R. Cahoon, J. Donnelly, M. Kearney, A.S. Kolker, L.L. Leonard, R.A. Orson, and J.C. Stevenson, 2008: “Site-Specific Scenarios for Wetlands Accretion in the Mid-Atlantic Region. Section 2.1” in *Background Documents Supporting Climate Change Science Program Synthesis and Assessment Product 4.1: Coastal Elevations and Sensitivity to Sea Level Rise*, J.G. Titus and E.M. Strange (eds.), EPA430R07004, Washington, DC: U.S. EPA.  
[http://www.epa.gov/climatechange/effects/downloads/section2\\_1.pdf](http://www.epa.gov/climatechange/effects/downloads/section2_1.pdf)
- Stevenson and Kearney, 2008, “Impacts of Global Climate Change and Sea-Level Rise on Tidal Wetlands” Pending chapter of manuscript by University of California Press.
- Titus, J.G., R.A. Park, S.P. Leatherman, J.R. Weggel, M.S. Greene, P.W. Mausel, M.S. Trehan, S. Brown, C. Grant, and G.W. Yohe. 1991. Greenhouse Effect and Sea Level Rise: Loss of Land and the Cost of Holding Back the Sea. *Coastal Management* 19:2:171-204.
- United States Fish and Wildlife Service, Federal Highway Administration Western Federal Lands Highway Division. 2009. Environmental Assessment for the Ni-les’tun Unit of the Pelican Island National Wildlife Refuge Restoration and North Bank Land Improvement Project

## Appendix A: Contextual Results


The SLAMM model does take into account the context of the surrounding lands or open water when calculating effects. For example, erosion rates are calculated based on the maximum fetch (wave action) which is estimated by assessing contiguous open water to a given marsh cell. Another example is that inundated dry lands will convert to marshes or ocean beach depending on their proximity to open ocean.

For this reason, an area larger than the boundaries of the USFWS refuge was modeled. These results maps are presented here with the following caveats:


- Results were closely examined (quality assurance) within USFWS refuges but not closely examined for the larger region.
- Site-specific parameters for the model were derived for USFWS refuges whenever possible and may not be regionally applicable.
- Especially in areas where dikes are present, an effort was made to assess the probable location and effects of dikes for USFWS refuges, but this effort was not made for surrounding areas.




Pelican Island National Wildlife Refuge within simulation context (black).




Pelican Island Context, Initial Condition




Pelican Island Context, 2025, Scenario A1B Mean




Pelican Island Context, 2050, Scenario A1B Mean




Pelican Island Context, 2075, Scenario A1B Mean




Pelican Island Context, 2100, Scenario A1B Mean




Pelican Island Context, Initial Condition




Pelican Island Context, 2025, Scenario A1B Maximum




Pelican Island Context, 2050, Scenario A1B Maximum




Pelican Island Context, 2075, Scenario A1B Maximum




Pelican Island Context, 2100, Scenario A1B Maximum




Pelican Island Context, Initial Condition




Pelican Island Context, 2025, 1 meter




Pelican Island Context, 2050, 1 meter




Pelican Island Context, 2075, 1 meter




Pelican Island Context, 2100, 1 meter




Pelican Island Context, Initial Condition




Pelican Island Context, 2025, 1.5 meter




Pelican Island Context, 2050, 1.5 meter




Pelican Island Context, 2075, 1.5 meter




Pelican Island Context, 2100, 1.5 meter




Pelican Island Context, Initial Condition




Pelican Island Context, 2025, 2 meter



Pelican Island Context, 2050, 2 meter



Pelican Island Context, 2075, 2 meter



Pelican Island Context, 2100, 2 meter