

# Application of the Sea-Level Affecting Marshes Model (SLAMM 5.0) to Sandy Point National Wildlife Refuge

Prepared For: Dr. Brian Czech, Conservation Biologist

U. S. Fish and Wildlife Service  
National Wildlife Refuge System  
Division of Natural Resources and Conservation Planning  
Conservation Biology Program  
4401 N. Fairfax Drive - MS 670  
Arlington, VA 22203

July 10, 2008

Jonathan S. Clough, Warren Pinnacle Consulting, Inc.  
PO Box 253, Warren VT, 05674  
(802)-496-3476

# Application of the Sea-Level Affecting Marshes Model (SLAMM 5.0) to Sandy Point National Wildlife Refuge

<b>Introduction.....</b>	<b>1</b>
<b>Model Summary .....</b>	<b>1</b>
Sea-Level Rise Scenarios .....	2
<b>Methods and Data Sources .....</b>	<b>4</b>
<b>Results .....</b>	<b>8</b>
<b>Discussion: .....</b>	<b>13</b>
<b>References .....</b>	<b>14</b>
<b>Appendix A: Contextual Results .....</b>	<b>16</b>

## Introduction

Tidal marshes are among the most susceptible ecosystems to climate change, especially accelerated sea level rise (SLR). Sea level is predicted to increase by 30 cm to 100 cm by 2100 based on the International Panel on Climate Change (IPCC) Special Report on Emissions Scenarios (SRES) (Meehl et al. 2007). Rising sea level may result in tidal marsh submergence (Moorhead and Brinson 1995) and habitat migration as salt marshes transgress landward and replace tidal freshwater and brackish marsh (Park et al. 1991).

In an effort to address the potential effects of sea level rise on United States national wildlife refuges, the U. S. Fish and Wildlife Service contracted the application of the SLAMM model for most Region 4 refuges. This analysis is designed to assist in the production of comprehensive conservation plans (CCPs) for each refuge. A CCP is a document that provides a framework for guiding refuge management decisions. All refuges are required by law to complete a CCP by 2012.

## Model Summary

Changes in tidal marsh area and habitat type in response to sea-level rise were modeled using the Sea Level Affecting Marshes Model (SLAMM 5.0) that accounts for the dominant processes involved in wetland conversion and shoreline modifications during long-term sea level rise (Park et al. 1989; [www.warrenpinnacle.com/prof/SLAMM](http://www.warrenpinnacle.com/prof/SLAMM)).

Successive versions of the model have been used to estimate the impacts of sea level rise on the coasts of the U.S. (Titus et al., 1991; Lee, J.K., R.A. Park, and P.W. Mause. 1992; Park, R.A., J.K. Lee, and D. Canning 1993; Galbraith, H., R. Jones, R.A. Park, J.S. Clough, S. Herrod-Julius, B. Harrington, and G. Page. 2002; National Wildlife Federation et al., 2006; Glick, Clough, et al. 2007; Craft et al., 2009).

Within SLAMM, there are five primary processes that affect wetland fate under different scenarios of sea-level rise:

- **Inundation:** The rise of water levels and the salt boundary are tracked by reducing elevations of each cell as sea levels rise, thus keeping mean tide level (MTL) constant at zero. The effects on each cell are calculated based on the minimum elevation and slope of that cell.
- **Erosion:** Erosion is triggered based on a threshold of maximum fetch and the proximity of the marsh to estuarine water or open ocean. When these conditions are met, horizontal erosion occurs at a rate based on site-specific data.
- **Overwash:** Barrier islands of under 500 meters width are assumed to undergo overwash during each 25-year time-step due to storms. Beach migration and transport of sediments are calculated.
- **Saturation:** Coastal swamps and fresh marshes can migrate onto adjacent uplands as a response of the fresh water table to rising sea level close to the coast.

- **Accretion:** Sea level rise is offset by sedimentation and vertical accretion using average or site-specific values for each wetland category. Accretion rates may be spatially variable within a given model domain.

SLAMM Version 5.0 is the latest version of the SLAMM Model, developed in 2006/2007 and based on SLAMM 4.0. SLAMM 5.0 provides the following refinements:

- The capability to simulate fixed levels of sea-level rise by 2100 in case IPCC estimates of sea-level rise prove to be too conservative;
- Additional model categories such as “Inland Shore,” “Irregularly Flooded (Brackish) Marsh,” and “Tidal Swamp.”
- *Optional.* In a defined estuary, salt marsh, brackish marsh, and tidal fresh marsh can migrate based on changes in salinity, using a simple though geographically-realistic salt wedge model. This optional model was not used when creating results for Sandy Point.

Model results presented in this report were produced using SLAMM version 5.0.1 which was released in early 2008 based on only minor refinements to the original SLAMM 5.0 model. Specifically, the accretion rates for swamps were modified based on additional literature review. For a thorough accounting of SLAMM model processes and the underlying assumptions and equations, please see the SLAMM 5.0.1 technical documentation (Clough and Park, 2008). This document is available at <http://warrenpinnacle.com/prof/SLAMM>


## ***Sea-Level Rise Scenarios***

The primary set of eustatic (global) sea level rise scenarios used within SLAMM was derived from the work of the Intergovernmental Panel on Climate Change (IPCC 2001). SLAMM 5 was run using the following IPCC and fixed-rate scenarios:

<b>Scenario</b>	<b>Eustatic SLR by 2025 (cm)</b>	<b>Eustatic SLR by 2050 (cm)</b>	<b>Eustatic SLR by 2075 (cm)</b>	<b>Eustatic SLR by 2100 (cm)</b>
A1B Mean	8	17	28	39
A1B Max	14	30	49	69
1 meter	13	28	48	100
1.5 meter	18	41	70	150

Recent literature (Chen et al., 2006, Monaghan et al., 2006) indicates that the eustatic rise in sea levels is progressing more rapidly than was previously assumed, perhaps due to the dynamic changes in ice flow omitted within the IPCC report’s calculations. A recent paper in the journal *Science* (Rahmstorf, 2007) suggests that, taking into account possible model error, a feasible range by 2100 might be 50 to 140 cm. To allow for flexibility when interpreting the results, SLAMM was also run assuming 1 meter, 1½ meters of eustatic sea-level rise by the year 2100. The A1B- maximum scenario was scaled up to produce these bounding scenarios (Figure 1).

**Figure 1: Summary of SLR Scenarios Utilized**



## Methods and Data Sources


No LIDAR data were found for the Sandy Point NWR so elevation data for Sandy Point are based on the National Elevation Dataset (NED). An examination of the metadata for the NED indicates that it was derived from the 1954 USGS topographic map shown below (Figure 2).

**Figure 2: Sandy Point from USGS Map.**  
Dotted line is 10 foot contour other lines are 20 foot contours.



There are very few contours displayed indicating that there is considerable uncertainty where the elevation of the dry land is between zero and the ten foot contour.

**Figure 3: Sandy Point 3D elevation map as derived from the NED**



Based on the National Elevation Dataset there is one significant plateau at roughly three meters above sea level.

Another available elevation data set is the Shuttle Radar Topography Mission (SRTM). (<http://www2.jpl.nasa.gov/srtm/>) This data set does not have a high degree of vertical resolution for this type of modeling purpose, however (the 90% absolute error for vertical resolution is 6.5 meters, Rodriguez et. al 2005). An examination of the SRTM data set was undertaken but it provided similar results of a plateau that is roughly 3 meters above mean tide level. For this reason, the NED elevation data was utilized as the best available data-set.

The National Wetlands Inventory for the Sandy Point region suggests that it is exclusively composed of dry land. No other wetlands categories or ocean beach categories are indicated in that data set. However, photos of sandy point and examination of satellite photos indicate that there is roughly a 30 meter fringe of ocean beach. Moreover, this beach is a critical habitat for leatherback turtles. Using satellite photographs a 30 meter fringe of beach was added to the NWI data-set and beach elevations were set using the SLAMM elevation pre-processor (which produces a uniform distribution of beach elevations starting at mean low water and progressing to mean high water spring). This process is obviously subject to considerable uncertainty but seems closer to reality than modeling “Sandy Point” with no beach at all.



Conversations with Michael Evans, Refuge Manager and Dr. Marc Albrecht out of University of Nebraska were undertaken to try to find better data sets for elevation and to get a sense for the site. Michael Evans suggested that the beach very “mobile,” that is to say that it is constantly eroding and reconfiguring itself based on wave action. This type of cyclical erosion and recreation of beaches is not something that can be predicted using the SLAMM model and projections of such dynamic action would likely be subject to considerable uncertainty using any model.

The historic trend for Sea Level Rise was calculated at 0.965 mm/year based on long term trends measured at Charlotte Amalie, Virgin Islands (NOAA station 9751639) and San Juan, Puerto Rico (9755371). This historical trend is lower than the global average that is approximately 1.5 mm/year indicating that local effects in this region protect it to some degree from the effects of sea level rise.

There is significant uncertainty relating the elevation data which is listed as having a vertical datum of NAVD88 to Mean Sea Level for the Caribbean Region. The NAVD88 correction is generally available through the NOAA CO-OPS database but not for the Caribbean region, possibly because NAVD88 is not considered “relevant” to the Caribbean. Nonetheless, NED metadata indicates that elevation data from the Caribbean are related to the NAVD88 basis.

The estimate for this relationship used in modeling was derived using the average of the closest NOAA stations in the CO-OPS database being Ocean Reef Harbor, Key Largo (#8723519); Rock Harbor Inside (#8723688); Duck Key (#8723927); Key Colony Beach (#8723962), and Islamorada, Whale Harbor Channel (#8723797). The average correction was -0.28 meters. This was the same procedure used in (Padilla Nieves, 2008) when the model was applied to Puerto Rico. That paper states: “transferring such leveling data carries significant uncertainty because NAVD88 elevation is based on site specific mean sea levels that may not be consistent from one location to another.”

The oceanic tide range was calculated using the average of the two closest NOAA stations, being: St. Johns Island, Coral Harbor, VI (9751373) and Lime Tree Bay, St. Croix, VI (9751401).

Parameters pertaining to marshes (i.e. accretion rates and erosion rates) are not relevant to this site as there are no wetlands identified based on the National Wetlands Inventory, nor are any predicted to occur. Default values are therefore used, though the model will not be sensitive to those choices.

Modeled U.S. Fish and Wildlife Service refuge boundaries are based on Approved Acquisition Boundaries as received from Kimberly Eldridge, lead cartographer with U.S. Fish and Wildlife Service, and are current as of June, 2008.

The cell-size used for this analysis was 30 meter by 30 meter cells. However, the SLAMM model does track partial conversion of cells based on elevation and slope.



## SLAMM INPUT PARAMETERS FOR Sandy Point

Site	Sandy Point
NED Source Date (yyyy)	1954
NWI_photo_date (yyyy)	1985
Direction_OffShore (N S E W)	W
Historic_trend (mm/yr)	0.965
NAVD88_correction (MTL-NAVD88 in meters)	-0.28
Water Depth (m below MLW- N/A)	2
TideRangeOcean (meters: MHHW-MLLW)	0.216
TideRangeInland (meters)	0.216
Mean High Water Spring (m above MTL)	0.144
MHSW Inland (m above MTL)	0.144
Marsh Erosion (horz meters/year)	1.8
Swamp Erosion (horz meters/year)	1
TFlat Erosion (horz meters/year) [from 0.5]	2
Salt marsh vertical accretion (mm/yr) Final	3.9
Brackish March vert. accretion (mm/yr) Final	4.7
Tidal Fresh vertical accretion (mm/yr) Final	5.9
Beach/T.Flat Sedimentation Rate (mm/yr)	0.5
Frequency of Large Storms (yr/washover)	25

## Results

Predicted effects on dry land are insignificant until the sea level rise scenario extends to 1.5 meters at which point 2.5% of dry lands are predicted lost. These predictions are highly uncertain given the 10 foot vertical resolution of the input data. These predictions should be revisited if and when a high quality elevation data set is produced for the island.

The vast majority of ocean beach is predicted to be lost under all scenarios run. Again, though, this estimate is subject to significant uncertainty because ocean beach was not actually present in the national wetlands inventory and had to be manually added with elevations estimated based on the tide range. Within SLAMM, dry lands can convert to ocean beach as well, when they are inundated. However, uncertainty in the dry land elevations make it unlikely that this conversion is being precisely estimated.

Maps of SLAMM input and output to follow will use the following legend:




Sandy Point


IPCC Scenario A1B-Mean, 0.39 M SLR Eustatic by 2100

Results in Acres


	Initial	2025	2050	2075	2100
Dry Land	409.9	409.5	409.0	408.7	408.4
Estuarine Open Water	99.4	102.6	107.7	110.1	110.1
Open Ocean	28.9	33.0	38.8	40.0	40.0
Estuarine Beach	11.1	7.1	2.3	0.0	0.0
Ocean Beach	11.1	7.2	1.7	0.7	0.9
Inland Open Water	4.0	4.0	4.0	4.0	4.0
Tidal Flat	0.0	0.8	0.4	0.4	0.4
Trans. Salt Marsh	0.0	0.1	0.4	0.4	0.4
Saltmarsh	0.0	0.0	0.0	0.1	0.2
<b>Total (incl. water)</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>




Sandy Point Initial Condition




Sandy Point 2075 IPCC Scenario A1B-Mean



Sandy Point 2025 IPCC Scenario A1B-Mean



Sandy Point 2100 IPCC Scenario A1B-Mean




Sandy Point 2050 IPCC Scenario A1B-Mean

**Sandy Point**


IPCC Scenario A1B-Max, 0.69 M SLR Eustatic by 2100

Results in Acres


	Initial	2025	2050	2075	2100
Dry Land	409.9	409.3	408.9	408.5	407.5
Estuarine Open Water	99.4	107.0	110.1	110.4	111.3
Open Ocean	28.9	37.1	40.2	40.5	40.7
Estuarine Beach	11.1	3.0	0.0	0.0	0.0
Ocean Beach	11.1	3.2	0.3	0.2	0.6
Inland Open Water	4.0	4.0	4.0	4.0	4.0
Tidal Flat	0.0	0.5	0.4	0.4	0.0
Trans. Salt Marsh	0.0	0.3	0.3	0.1	0.2
Saltmarsh	0.0	0.0	0.3	0.3	0.1
<b>Total (incl. water)</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>




Sandy Point Initial Condition




Sandy Point 2075 IPCC Scenario A1B-Max.



Sandy Point 2025 IPCC Scenario A1B-Max.



Sandy Point 2100 IPCC Scenario A1B-Max.




Sandy Point 2050 IPCC Scenario A1B-Max.


Sandy Point  
1 Meter Eustatic SLR by 2100

Results in Acres


	Initial	2025	2050	2075	2100
Dry Land	409.9	409.2	409.0	407.8	405.7
Estuarine Open Water	99.4	110.3	110.6	111.3	111.9
Open Ocean	28.9	40.0	40.3	40.3	41.1
Estuarine Beach	11.1	0.1	0.0	0.0	0.0
Ocean Beach	11.1	0.2	0.0	0.7	0.7
Inland Open Water	4.0	4.0	4.0	4.0	4.0
Tidal Flat	0.0	0.0	0.0	0.0	0.0
Trans. Salt Marsh	0.0	0.5	0.1	0.2	0.7
Saltmarsh	0.0	0.2	0.5	0.1	0.2
<b>Total (incl. water)</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>




Sandy Point Initial Condition




Sandy Point 2075 1 meter Eustatic by 2100



Sandy Point 2025 1 meter Eustatic by 2100



Sandy Point 2100 1 meter Eustatic by 2100




Sandy Point 2050 1 meter Eustatic by 2100

Sandy Point


1.5 Meters Eustatic SLR by 2100

Results in Acres


	Initial	2025	2050	2075	2100
Dry Land	409.9	409.3	408.5	405.4	399.5
Estuarine Open Water	99.4	110.5	110.8	112.0	113.3
Open Ocean	28.9	40.0	40.0	41.1	42.8
Estuarine Beach	11.1	0.0	0.0	0.0	0.0
Ocean Beach	11.1	0.0	0.3	0.9	1.1
Inland Open Water	4.0	4.0	4.0	4.0	4.0
Tidal Flat	0.0	0.0	0.0	0.0	0.0
Trans. Salt Marsh	0.0	0.5	0.2	0.8	3.0
Saltmarsh	0.0	0.0	0.5	0.2	0.8
<b>Total (incl. water)</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>	<b>564.4</b>




Sandy Point Initial Condition




Sandy Point 2075 1.5 meter Eustatic by 2100



Sandy Point 2025 1.5 meter Eustatic by 2100



Sandy Point 2100 1.5 meter Eustatic by 2100



Sandy Point 2050 1.5 meter Eustatic by 2100

## Discussion:

SLAMM results for the Sandy Point NWR are subject to considerable uncertainty due to limitations in both the national elevation dataset and the national wetlands inventory. Uncertainty with respect to the loss of dry land could be reduced if a higher resolution elevation data set (with higher vertical precision) were created or located.

It may be especially difficult to successfully model the fate of the ocean beach at Sandy Point given this type of a “mobile” beach in which the beach regularly erodes and is regenerated (Personal communication with U.S. FWS refuge manager Michael Evans.) The SLAMM model beach-loss results presented here utilize the Bruun rule to estimate beach erosion and estimate inundation based on estimated beach elevations. As there was no beach cover-class indicated in the National Wetlands Inventory, this SLAMM beach-loss analysis is especially uncertain, even with respect to initial conditions.

Some ocean beach may be created due to the inundation of dry lands, but the predicted extent of this inundation and conversion to beach-lands is subject to the quality of the elevation data set for the dry lands.



## References

- Cahoon, D.R., J. W. Day, Jr., and D. J. Reed, 1999. "The influence of surface and shallow subsurface soil processes on wetland elevation: A synthesis." *Current Topics in Wetland Biogeochemistry*, 3, 72-88.
- Chen, J. L., Wilson, C. R., Tapley, B. D., 2006 "Satellite Gravity Measurements Confirm Accelerated Melting of Greenland Ice Sheet" *Science* 2006 0: 1129007
- Clough, J.S. and R.A. Park, 2007, *Technical Documentation for SLAMM 5.0.1* February 2008, Jonathan S. Clough, Warren Pinnacle Consulting, Inc, Richard A. Park, Eco Modeling.  
<http://warrenpinnacle.com/prof/SLAMM>
- Craft C, Clough J, Ehman J, Guo H, Joye S, Machmuller M, Park R, and Pennings S. Effects of Accelerated Sea Level Rise on Delivery of Ecosystem Services Provided by Tidal Marshes: A Simulation of the Georgia (USA) Coast. *Frontiers in Ecology and the Environment*. 2009; 7, doi:10.1890/070219
- Galbraith, H., R. Jones, R.A. Park, J.S. Clough, S. Herrod-Julius, B. Harrington, and G. Page. 2002. Global Climate Change and Sea Level Rise: Potential Losses of Intertidal Habitat for Shorebirds. *Waterbirds* 25:173-183.
- Glick, Clough, et al. *Sea-level Rise and Coastal Habitats in the Pacific Northwest An Analysis for Puget Sound, Southwestern Washington, and Northwestern Oregon* July 2007  
<http://www.nwf.org/sealevelrise/pdfs/PacificNWSeaLevelRise.pdf>
- IPCC, 2001: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change* [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881pp.
- Lee, J.K., R.A. Park, and P.W. Mausel. 1992. Application of Geoprocessing and Simulation Modeling to Estimate Impacts of Sea Level Rise on the Northeast Coast of Florida. *Photogrammetric Engineering and Remote Sensing* 58:11:1579-1586.
- Meehl GA, Stocker TF, Collins WD, Friedlingstein P, Gaye AT, Gregory JM, Kitoh A, Knutti R, Murphy JM, Noda A, Raper SCB, Watterson IG, Weaver AJ and Zhao ZC. 2007. Global climate projections. Pp. 747-845. In: Solomon S, Qin, D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor, M and Miller HL, (eds.) *Climate change 2007: The physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK: Cambridge University Press.
- Monaghan, A. J. *et al*, 2006 "Insignificant Change in Antarctic Snowfall Since the International Geophysical Year" *Science* 2006 313: 827-831.
- Moorhead, KK and Brinson MM. 1995. Response of wetlands to rising sea level in the lower coastal plain of North Carolina. *Ecological Applications* 5: 261-271.

- National Wildlife Federation et al., *An Unfavorable Tide: Global Warming, Coastal Habitats and Sportfishing in Florida* 4, 6 (2006).  
<http://www.targetglobalwarming.org/files/AnUnfavorableTideReport.pdf>
- Park, R.A., J.K. Lee, and D. Canning. 1993. Potential Effects of Sea Level Rise on Puget Sound Wetlands. *Geocarto International* 8(4):99-110.
- Park, R.A., M.S. Trehan, P.W. Mause, and R.C. Howe. 1989a. The Effects of Sea Level Rise on U.S. Coastal Wetlands. In *The Potential Effects of Global Climate Change on the United States: Appendix B - Sea Level Rise*, edited by J.B. Smith and D.A. Tirpak, 1-1 to 1-55. EPA-230-05-89-052. Washington, D.C.: U.S. Environmental Protection Agency.
- Rahmstorf, Stefan 2007, "A Semi-Empirical Approach to Projecting Future Sea-Level Rise," *Science* 2007 315: 368-370.
- Rodriguez, E., C.S. Morris, J.E. Belz, E.C. Chapin, J.M. Martin, W. Daffer, S. Hensley, 2005, *An assessment of the SRTM topographic products*, Technical Report JPL D-31639, Jet Propulsion Laboratory, Pasadena, California, 143 pp.
- Reed, D.J., D.A. Bishara, D.R. Cahoon, J. Donnelly, M. Kearney, A.S. Kolker, L.L. Leonard, R.A. Orson, and J.C. Stevenson, 2008: "Site-Specific Scenarios for Wetlands Accretion in the Mid-Atlantic Region. Section 2.1" in *Background Documents Supporting Climate Change Science Program Synthesis and Assessment Product 4.1: Coastal Elevations and Sensitivity to Sea Level Rise*, J.G. Titus and E.M. Strange (eds.), EPA430R07004, Washington, DC: U.S. EPA.  
[http://www.epa.gov/climatechange/effects/downloads/section2\\_1.pdf](http://www.epa.gov/climatechange/effects/downloads/section2_1.pdf)
- Stevenson and Kearney, 2008, "Impacts of Global Climate Change and Sea-Level Rise on Tidal Wetlands" Pending chapter of manuscript by University of California Press.
- Titus, J.G., R.A. Park, S.P. Leatherman, J.R. Weggel, M.S. Greene, P.W. Mause, M.S. Trehan, S. Brown, C. Grant, and G.W. Yohe. 1991. Greenhouse Effect and Sea Level Rise: Loss of Land and the Cost of Holding Back the Sea. *Coastal Management* 19:2:171-204.

## Appendix A: Contextual Results


The SLAMM model does take into account the context of the surrounding lands or open water when calculating effects. For example, erosion rates are calculated based on the maximum fetch (wave action) which is estimated by assessing contiguous open water to a given marsh cell. Another example is that inundated dry lands will convert to marshes or ocean beach depending on their proximity to open ocean.

For this reason, an area larger than the boundaries of the USFWS refuge was modeled. These results maps are presented here with the following caveats:


- Results were closely examined (quality assurance) within USFWS refuges but not closely examined for the larger region.
- Site-specific parameters for the model were derived for USFWS refuges whenever possible and may not be regionally applicable.
- Especially in areas where dikes are present, an effort was made to assess the probable location and effects of dikes for USFWS refuges, but this effort was not made for surrounding areas.




Croix Initial Condition IPCC Scenario A1B-Mean




Croix 2025 IPCC Scenario A1B-Mean




Croix 2050 IPCC Scenario A1B-Mean




Croix 2075 IPCC Scenario A1B-Mean




Croix 2100 IPCC Scenario A1B-Mean



Croix Initial Condition




Croix 2025 IPCC Scenario A1B-Maximum




Croix 2050 IPCC Scenario A1B-Maximum




Croix 2075 IPCC Scenario A1B-Maximum




Croix 2100 IPCC Scenario A1B-Maximum



Croix Initial Condition




Croix 2025 1 meter Eustatic by 2100




Croix 2050 1 meter Eustatic by 2100







Croix 2075 1 meter Eustatic by 2100




Croix 2100 1 meter Eustatic by 2100




Croix Initial Condition




Croix 2025 1.5 meter Eustatic by 2100



Croix 2050 1.5 meter Eustatic by 2100



Croix 2075 1.5 meter Eustatic by 2100



Croix 2100 1.5 meter Eustatic by 2100